

Acute Chest Syndrome in Sickle Cell Disease Care Guideline



Inclusion Criteria:

- Children with sickle cell disease
- New pulmonary infiltrate on chest x-ray, involving at least one complete lung segment, excluding atelectasis

Assessment

- Admit to 5S or PICU
- NPO
- History and physical with focus on pulmonary exam
- Vitals, accurate height and weight
- Strict I/O; Daily Weight
- O2 saturation
- Diagnostics: CBC with Retic, CMP with LDH, STAT Type and Screen, Hgb Electrophoresis
- Chest X-ray (CXR) – 2 View
- For significant respiratory distress, consider VBG
- Fever >38°C or elevated WBC, obtain Blood Culture
- URI symptoms, obtain Respiratory Panel PCR



Interventions

- Stat Hematology consult (if not admitted to Hematology service)
- Pulmonary consult
- Notify apheresis team as soon as possible during working hours if anticipated procedure
- Consult PICU for possible apheresis catheter placement
- Oxygen to keep O2 sat $\geq 94\%$
- Maintenance IV fluids
- Antibiotics: cefTRIAxone 75 mg/kg IV q24h (max dose 2g) and azithromycin 10 mg/kg IV q24h (max dose 500mg)
- Pain medication: Ketorolac 0.5 mg/kg IV q6h (max dose 30mg) for 5 DAYS PRN Pain (Moderate)
- Cardiac monitoring
- Simple transfusion of PRBCs to **max** Hgb = 10 gm/dl
- Repeat CXR, CBC STAT if worsening respiratory symptoms/increased oxygen requirements
- Consider exchange transfusion (see order set) if:
 - initial Hgb ≥ 10 gm/dl
 - worsening respiratory symptoms
 - worsening radiological findings (CXR), despite simple transfusion and supportive care
- Albuterol HFA MDI 2 PUFFS Q4hr PRN wheezing or poor air exchange; Chest Physiotherapy after each treatment - Use Vest if patient > 10 kg (See Policy RT205)
- Incentive Spirometry (IS) for patients ≥ 5 years, every 2 hrs while awake - use bubbles or pinwheel if patient < 5 years or unable to do IS

Further Recommendations

- Consider furosemide for s/s of fluid overload
- Consider systemic steroid if wheezing
- Cardiology evaluation with Echo and EKG
- Keep Hgb > 9 gm/dl for 2-3 months post discharge

Discharge Criteria

- Afebrile > 24 hours
- Baseline oxygen requirement; improved respiratory symptoms
- On oral antibiotics
- Stable Hgb
- Adequate pain control on oral medication
- Adequate oral intake
- Catch up vaccines, specifically Pneumovax 23 (every 5 yrs); seasonal influenza; COVID-19 (recommended, if age appropriate)

Recommendations/ Considerations

Predictors:

- Pain crisis involving chest, shoulders and back
- Post anesthesia complication
- Respiratory Infections
- On narcotics
- Asthma exacerbation
- Baseline Hgb level may run low < 9gm/dl
- If suspected pulmonary embolism, obtain CT angiogram of chest
- May need more than 1 exchange transfusion if clinical findings not improving
- After recovery from acute crisis, patient should be started on hydroxyurea if not already taking; optimize dose
- History of more than 1 acute chest crisis, consider chronic transfusion protocols to keep HgbS < 25%
- If recurrent crises, consider BMT if match available
- Monitor for right ventricular (RV) failure and/or pulmonary hypertension
- If continued worsening despite above interventions, will need respiratory support, including non-invasive ventilation, intubation, iNO, or ECMO
- Apheresis team hours 7a-7p weekdays, 7a-7p weekends (see call schedule on PAWS)

Patient/Parent or Caregiver Education

- Asthma education/AAP if has evidence of asthma

References

- Crabtree, E. A., Mariscalco, M. M., Hesselgrave, J., Iniguez, S. F., Hilliard, T. J., Katkin, J. P., . . . Hockenberry, M. J. (2011). Improving care for children with sickle cell disease/acute chest syndrome. *Pediatrics*, *127*(2), e480-e488. <https://doi:10.1542/peds.2010-3099> (**Level III**)
- Howard, J., Hart, N., Roberts-Harewood, M., Cummins, M., Awogbade, M., & Davis, B. (2015). Guideline on the management of acute chest syndrome in sickle cell disease. *British Journal of Haematology*, *169*, 492-505. <https://doi:10.1111/bjh.13348> (**Level III**)
- Miller, S. T. (2011). How I treat acute chest syndrome in children with sickle cell disease. *Blood*, *117*, 5297-5305. <https://doi:10.1182/blood-2010-11-261834> (**Level III**)
- Texas Children's Hospital. (2008). Evidence-Based Clinical Decision Support Acute Chest Syndrome (ACS) Guideline.
- U.S. Department of Health and Human Services, National Institutes of Health. (2014). Evidence-Based Management of Sickle Cell Disease. *Expert Panel Report*. (**Level V**)
- Vichinsky, E. P., Neumayr, L. D., Earles, A. N., Williams, R., Lennette, E. T., Dean, D., . . . Manci, E. A. (2000). Causes and outcomes of the acute chest syndrome in sickle cell disease. National Acute Chest Syndrome Study Group. *The New England Journal of Medicine*, *342*(25), 1855-1865. <https://doi:10.1056/NEJM200006223422502> (**Level III**)