

Blunt abdominal trauma with any of the following¹:

<ul style="list-style-type: none"> GCS \leq 13 Abdominal bruising Abdominal Seatbelt Sign Abdominal distention Abdominal pain/tenderness Vomiting 	<ul style="list-style-type: none"> Abnormal FAST Abnormal chest or pelvic x-ray ALT >125 U/L; AST >200 U/L Decreased Hemoglobin (less than 7g/dl) Elevated Lipase Hematuria (UA >50 rbc/hpf)
--	---

Exclusion criteria: Blunt Abdominal Trauma with peritonitis and penetrating abdominal trauma

Manage other injuries

Unstable

Circulatory Status
(recent/ongoing bleeding suspected?)

Stable

CT abd/pelvis with IV contrast

Fluid Resuscitation
20 ml/kg NS or LR

Sustained response to bolus?

Resuscitation with PRBCs 10-20ml/kg

Recurrent hypotension or persistent shock^{2ab}

Negative

- No evidence of Hollow or solid organ injury
- If intraperitoneal free fluid:
 - Small volume isolated intraperitoneal free fluid with \leq 25 HU⁴

- Suspected Hollow organ injury
- >25 HU⁴ intraperitoneal free fluid
- Free air
- Soft tissue seatbelt sign
- Bowel wall thickening
- Mesenteric hematoma or stranding
- Abdominal Wall hernia

Significant intra-abdominal blood or contrast extravasation
OR
Solid organ injury

Other
Manage per surgeon discretion

Discharge Home from ED if:

- Vitals normal
- Tolerating diet
- Minimal abdominal pain
- No blood product administration in ED
- Access to care: transportation & phone

(admit to non-ICU for observation if all the above conditions are not met)

At discharge:

- Provide discharge instructions & return precautions.

See Hollow Viscus Injury Algorithm

See Blunt Liver/Spleen Injury

Failure of NOM

Angiography & Embolization³

Surgery³

NOM at Surgeon's discretion

1-Based on PECARN & PedSRC prediction rule
2a- Consider other causes (head injury, tension pneumothorax, tamponade, pelvic hemorrhage)
2b-Activate Massive Transfusion protocol (see MTP guideline)
3- must be available within 30 minutes of notification
4 – Hounsfield Units

Guideline adapted from the ATOMAC Blunt Pediatric Liver/Spleen Injury Guideline v12.

Clinical Care Guideline

Name: Blunt Abdominal Trauma
Emergency Department, Inpatient and ICU

I. Inclusion Criteria	<ul style="list-style-type: none"> • Patients ages 0-17 years presenting to CHOC Emergency Department with blunt abdominal trauma.
II. Exclusion Criteria	<ul style="list-style-type: none"> • Major multiple trauma when the blunt abdominal trauma does not direct the patient's plan of care. • Specific findings/Injuries: <ul style="list-style-type: none"> ○ Peritonitis ○ Pancreatic injuries ○ Penetrating injury • Significant medical comorbidities
III. Assessment and evaluation	<ul style="list-style-type: none"> • Review of history, physical exam, labs, and imaging. • Refer to Blunt Abdominal Trauma algorithm (attached) for recommendations on when to obtain CT abdomen/pelvis imaging and evaluation of the trauma patient with confirmed blunt abdominal trauma without peritonitis.
IV. Interventions (utilizing GRADE whenever possible)	<ul style="list-style-type: none"> • 1A Management of the awake and oriented blunt abdominal trauma patient starts with the primary survey, E-FAST, physical examination and the secondary survey, blood chemistry, vital signs followed by contrast-enhanced abdominal CT. • 1A CT is recommended for the evaluation of hemodynamically stable patients with equivocal findings on physical examination, associated neurologic injury, or multiple extra-abdominal injuries. Under these circumstances, patients with a negative CT should be admitted for observation. • 1A CT is the diagnostic modality of choice for nonoperative management of solid visceral injuries. • 2C Patients not admitted to the ICU and without signs of ongoing bleeding may be allowed to drink and eat when comfortable and able. • 3A: Objective diagnostic testing (i.e. CT) is indicated for patient with abnormal mentation, equivocal findings on physical examination, multiple injuries, concomitant chest injury or hematuria.
V. Recommendations and considerations (utilizing GRADE whenever possible)	<p>Considerations related to Operative Management</p> <ul style="list-style-type: none"> • 1A The presence of highly specific CT findings such as extraluminal air, extraluminal oral contrast, or bowel-wall defects warrants prompt surgical exploration. <p>Considerations related to Non-operative Management (NOM)</p>

- 1A NOM in the face of peritonitis should be excluded from this guideline and managed per physician discretion.
- 1A Failure to stabilize as noted by persistent or recurrent hypotension cannot have NOM dictated by algorithm alone without taking into account local resources and other injuries. These patients should be considered for surgery, urgent embolization, or continued NOM, depending on other injuries.
- 1B Patients at risk for missed injury, need for operation, or recurrent bleeding are those with multiple identified abdominal injuries (especially pancreas), those with a contrast blush on CT scan, and those with bicycle handlebar injuries, and may not be candidates for early discharge. Since delayed bleeds have occurred (often outside of the APSA guideline period), standardized education remains an important component of the discharge.
- 2A The presence of highly sensitive CT findings such as free fluid in the absence of solid organ injury, abnormal enhancement of bowel wall, and mesenteric stranding can be used as an adjunct to the clinical picture but should not solely determine management.

Considerations related to hemoglobin monitoring, ongoing bleeding and transfusion:

- 2C Clinical determination of recent or ongoing bleeding in children requires integration of multiple factors to determine the relative importance of SOI bleeding. Important factors to consider are listed on the algorithm instructions to assist in the determination recent bleeding significant enough to suggest shock.
- 2C Consider use of 1:1:1 transfusion ratios early in resuscitation.
- 2C Consideration for TEG-directed therapy may be given based on adult data.
- 2C Limiting crystalloid volume and early use of transfusion in children with significant bleeding should be considered based on the adult literature.
- 2C A pediatric practice management guideline may be used for all children under 18 years of age, but a caution about the use of the algorithm for children 9-16 years of age may be appropriate.
- 2D Transfusion of red blood cells in response to shock is recommended.

Considerations related to imaging:

- 1A Management of the awake and oriented blunt abdominal trauma patient starts with the primary survey, E-FAST, physical examination and the secondary survey, blood chemistry, vital signs followed by contrast-enhanced abdominal CT.
- 1A CT is recommended for the evaluation of hemodynamically stable patients with equivocal findings on physical examination, associated neurologic injury, or multiple extra-abdominal injuries. Under these

	<p>circumstances, patients with a negative CT should be admitted for observation.</p> <ul style="list-style-type: none"> • 1A CT is the diagnostic modality of choice for nonoperative management of solid visceral injuries. • 1A The presence of a seatbelt sign should prompt a CT scan and a high index of suspicion for bowel injury. • 2C Pediatric patients with Hounsfield units (HU) of 25 or less isolated intraperitoneal free fluid (IIF) and a non-peritonitic physical examination do not require operative exploration or further workup for intraabdominal injury. In the absence of other injuries, it is safe to discharge these patients without further workup. • 3A: Haemodynamically stable children with blunt trauma should undergo radiographic evaluation if they have gross haematuria or >50 red blood cells/high-power field on microscopic urine analysis. • 3: Low systolic blood pressure, abdominal tenderness, femur fracture, AST more than 200 U/L or ALT more than 125 U/L, an initial hematocrit less than 30% are important predictors of intra-abdominal trauma. <p>Considerations related to diagnosis and management:</p> <ul style="list-style-type: none"> • 1A In the context of blunt abdominal trauma with or without solid organ injury, bowel injuries are often missed. A high index of suspicion is required. • 2B Delay in the diagnosis of bowel injury is linked to increased morbidity and mortality. • 3: Elevated amylase/Lipase levels are suggestive but not diagnostic of pancreatic injury <p>Considerations related to discharge and follow-up:</p> <ul style="list-style-type: none"> • 2D Children may return to school when comfortable and able. • 2D Parents and patients are instructed to return to the emergency department for increasing pain, pallor, dizziness, difficulty breathing, vomiting, worsening shoulder pain, jaundice, gastrointestinal bleeding, or black tarry stools.
<p>Patient Family Education (PFE)</p>	<p>Lexicomp patient education:</p> <ul style="list-style-type: none"> • Blunt Abdominal Trauma (Appendix B)
<p>Discharge criteria</p>	<p>For all patients, they may be discharged when all of the below criteria are met:</p> <ol style="list-style-type: none"> 1. Vital signs are normal for age 2. Adequate oral intake 3. Minimal abdominal pain and/or pain controlled with oral medication

	4. Access to care: transportation & phone
Discharge instructions:	<ol style="list-style-type: none"> 1. Return to school: As tolerated 2. Follow-up: As needed 3. Return to the Emergency Department for increasing pain, pallor, dizziness, difficulty breathing, vomiting, worsening shoulder pain, jaundice, gastrointestinal bleeding, or black tarry stools.
References (leveled)	<p>Level II:</p> <p>Linnaus ME, Notrica DM, Langlais CS, St Peter SD, Leys CM, Ostlie DJ, Maxson RT, Ponsky T, Tuggle DW, Eubanks JW 3rd, Bhatia A, Alder AC, Greenwell C, Garcia NM, Lawson KA, Motghare P, Letton RW. Prospective validation of the shock index pediatric-adjusted (SIPA) in blunt liver and spleen trauma: An ATOMAC+ study. <i>J Pediatr Surg.</i> 2017 Feb;52(2):340-344. doi: 10.1016/j.jpedsurg.2016.09.060. Epub 2016 Sep 23. PMID: 27717564.</p> <p>Level III</p> <p>Borgialli, Dominic A et al. "Association between the seat belt sign and intra-abdominal injuries in children with blunt torso trauma in motor vehicle collisions." <i>Academic emergency medicine : official journal of the Society for Academic Emergency Medicine</i> vol. 21,11 (2014): 1240-8. doi:10.1111/acem.12506</p> <p>Gaffley, Michaela et al. "Evaluation of an evidence-based guideline to reduce CT use in the assessment of blunt pediatric abdominal trauma." <i>Journal of pediatric surgery</i> vol. 56,2 (2021): 297-301. doi:10.1016/j.jpedsurg.2020.07.002</p> <p>Holmes, James F et al. "Identification of children with intra-abdominal injuries after blunt trauma." <i>Annals of Emergency Medicine</i> vol. 39,5 (2002): 500-9. doi:10.1067/mem.2002.122900</p> <p>Holmes, James F et al. "Identifying children at very low risk of clinically important blunt abdominal injuries." <i>Annals of emergency medicine</i> vol. 62,2 (2013): 107-116.e2. doi:10.1016/j.annemergmed.2012.11.009</p> <p>Level IV</p>

Harrell, Kevin N et al. "Management of blunt traumatic abdominal wall hernias: A Western Trauma Association multicenter study." *The journal of trauma and acute care surgery* vol. 91,5 (2021): 834-840.
doi:10.1097/TA.0000000000003250

Magoteaux SR, Notrica DM, Langlais CS, Linnaus ME, Raines AR, Letton RW, Alder AC, Greenwell C, Eubanks JW, Lawson KA, Garcia NM, St Peter SD, Ostlie DJ, Leys CM, Bhatia A, Maxson RT, Tuggle DW, Ponsky TA. Hypotension and the need for transfusion in pediatric blunt spleen and liver injury: An ATOMAC+ prospective study. *J Pediatr Surg.* 2017 Jun;52(6):979-983. doi: 10.1016/j.jpedsurg.2017.03.021. Epub 2017 Mar 16. PMID: 28363471.

Level V

Adelgais, Kathleen M et al. "Accuracy of the abdominal examination for identifying children with blunt intra-abdominal injuries." *The Journal of pediatrics* vol. 165,6 (2014): 1230-1235.e5. doi:10.1016/j.jpeds.2014.08.014

Perea, Lindsey L et al. "Low-density Isolated Intraperitoneal Free Fluid in Pediatric Blunt Trauma Is Not Associated with Abdominal Injury." *Pediatric emergency care* vol. 38,1 (2022): e143-e146.
doi:10.1097/PEC.0000000000002189

Expert Opinion:

Notrica DM, Eubanks JW 3rd, Tuggle DW, Maxson RT, Letton RW, Garcia NM, Alder AC, Lawson KA, St Peter SD, Megison S, Garcia-Filion P. Nonoperative management of blunt liver and spleen injury in children: Evaluation of the ATOMAC guideline using GRADE. *J Trauma Acute Care Surg.* 2015 Oct;79(4):683-93. doi: 10.1097/TA.0000000000000808. PMID: 26402546.

Hoff, W., Holevar, M., Nagy, K., Patterson, L., Young, J., Arrillaga, A., Valenziano, C. (n.d.). Practice Management Guidelines for the Evaluation of Blunt Abdominal Trauma: The EAST Practice Management Guide

Santucci, R A et al. "Evaluation and management of renal injuries: consensus statement of the renal trauma subcommittee." *BJU*

	<p><i>international</i> vol. 93,7 (2004): 937-54. doi:10.1111/j.1464-4096.2004.04820.x</p> <p>Smyth, L., Bendinelli, C., Lee, N. <i>et al.</i> WSES guidelines on blunt and penetrating bowel injury: diagnosis, investigations, and treatment. <i>World J Emerg Surg</i> 17, 13 (2022). https://doi.org/10.1186/s13017-022-00418-y</p>
<p>Education and communication dissemination plans (provider, nursing, and other departments/ services as needed).</p>	<p>Guideline to be reviewed and distributed at the following:</p> <ol style="list-style-type: none"> 1. Trauma M&M and IOP 2. Trauma Nurse staff meetings 3. Critical Care Committee 4. ED Physician meetings
<p>Pertinent policies and/or procedures</p>	<ol style="list-style-type: none"> 1. Massive Blood Transfusion (F747)

Blunt Abdominal Trauma Discharge Instructions

Printed on 2024-02-23

You must carefully read the "Consumer Information Use and Disclaimer" below in order to understand and correctly use this information

About this topic

Trauma is an injury caused by some force or energy from the environment. Abdominal trauma can be very serious. Trauma calls for special doctors to care for you to help your chances of healing.

With abdominal trauma, you may have a mild injury like bruises. Other times you may have more serious problems if there is damage to any of your internal organs. You may need to have surgery right away. Trauma can lead to bleeding inside of your body, shock, or even death.

Most often, you will need care right away after a trauma. You may need emergency care where the accident happened. Staff will work to make sure you are breathing. They will control bleeding and protect you from any more harm. Then, you may be taken to the emergency room.

At the hospital, you will be given urgent medical care. The goal is to find and treat your injuries and to help you heal fully. How quickly you heal from a trauma may vary from person to person. Healing is based on how:

- Serious the injury and disability are
- Quickly treatment is given
- You respond to treatment

What care is needed at home?

- Ask your doctor what you need to do when you go home. Make sure you ask questions if you do not understand what the doctor says.
- Get lots of rest. You may have a problem sleeping or changes in sleep patterns. Be sure to take time to rest even if you cannot sleep. Take naps during the day.
- You may still have pain, even after fully healing. Talk with the doctor about your pain and how to control it.
- You may have emotional problems, worry, anxiety, or low mood. This could be caused by the trauma. The doctor may suggest counseling or other therapies.
- Talk to your doctor about how to care for your injuries. Ask your doctor about:
 - When you should change your bandages
 - When you may take a bath or shower
 - If you need to limit your activity or be on bedrest
 - If you need to be careful with lifting things over 10 pounds (4.5 kg)
 - When you may go back to your normal activities like work, driving, or sex
- Be sure to wash your hands before touching your wound or dressing.

What follow-up care is needed?

- Your condition needs close monitoring.
- Your doctor may ask you to make visits to the office to check on your progress. Be sure to keep these visits.

What drugs may be needed?

The doctor may order drugs to:

- Help with pain and swelling
- Prevent or fight an infection
- Prevent constipation

Will physical activity be limited?

You may need to rest for a while. You should not do physical activity that makes your health problem worse. If you run, work out, or play sports, you may not be able to do those things until your health problem gets better.

What problems could happen?

- Infection

- Bleeding
- Need for surgery

What can be done to prevent this health problem?

There are no specific ways to prevent general trauma. But, you can help prevent traumatic events by:

- Always wear a seat belt with a shoulder strap.
- Drive safely. Obey speed limits.
- Do not drink and drive. Do not text or talk on the phone when you drive.
- Do not use illegal drugs.
- Stay active and work out to keep your muscles and joints strong and flexible.
- Follow special precautions when in high-risk situations, like:
 - Avoid activities that may cause falls.
 - Wear protective gear when participating in sports.
- Talk to your doctor if you are worried about domestic abuse and be sure you have resources for your safety.

When do I need to call the doctor?

- Signs of infection. These include a fever of 100.4°F (38°C) or higher; chills, and redness near the site of injuries.
- Throwing up blood or passing blood with your stool
- Upset stomach and throwing up
- New blood in your urine or stool
- Increasing size of your belly
- Dizziness

Teach Back: Helping You Understand

The Teach Back Method helps you understand the information we are giving you. After you talk with the staff, tell them in your own words what you learned. This helps to make sure the staff has described each thing clearly. It also helps to explain things that may have been confusing. Before going home, make sure you are able to do these:

- I can tell you about my condition.
- I can tell you what changes I need to make with my activities.
- I can tell you what I will do if I have fever, throw up blood, or have an upset stomach.

Consumer Information Use and Disclaimer:

This generalized information is a limited summary of diagnosis, treatment, and/or medication information. It is not meant to be comprehensive and should be used as a tool to help the user understand and/or assess potential diagnostic and treatment options. It does NOT include all information about conditions, treatments, medications, side effects, or risks that may apply to a specific patient. It is not intended to be medical advice or a substitute for the medical advice, diagnosis, or treatment of a health care provider based on the health care provider's examination and assessment of a patient's specific and unique circumstances. Patients must speak with a health care provider for complete information about their health, medical questions, and treatment options, including any risks or benefits regarding use of medications. This information does not endorse any treatments or medications as safe, effective, or approved for treating a specific patient. UpToDate, Inc. and its affiliates disclaim any warranty or liability relating to this information or the use thereof. The use of this information is governed by the Terms of Use, available at <https://www.wolterskluwer.com/en/known-clinical-effectiveness-terms>

Last Reviewed Date

2021-03-29

 Wolters Kluwer

Copyright © 2024 UpToDate, Inc. and its affiliates and/or licensors. All rights reserved.